

Note on proton-antiproton suppression in 200 AGeV Au-Au collisions

Thorsten Renk* and Kari J. Eskola†

*Department of Physics, P.O. Box 35 FI-40014 University of Jyväskylä, Finland and
Helsinki Institute of Physics, P.O. Box 64 FI-00014, University of Helsinki, Finland*

We discuss the measured nuclear suppression of $p + \bar{p}$ production in 200 AGeV Au-Au collisions at RHIC within radiative energy loss. For the AKK set of fragmentation functions, proton production is dominated by gluons, giving rise to the expectation that the nuclear suppression for $p + \bar{p}$ should be stronger than for pions due to the stronger coupling of gluons to the quenching medium. Using a hydrodynamical description for the soft matter evolution, we show that this is indeed seen in the calculation. However, the expected suppression factors for pions and protons are sufficiently similar that a discrimination with present data is not possible. In the high p_T region above 6 GeV where the contributions of hydrodynamics and recombination to hadron production are negligible, the model calculation is in good agreement with the data on $p + \bar{p}$ suppression.

PACS numbers: 25.75.-q, 25.75.Gz

The STAR collaboration has measured the nuclear suppression ratio R_{CP} (yield for 0 – 12% central collisions divided by yield for 60 – 80% peripheral collisions) for $\pi^+ + \pi^-$ and $p + \bar{p}$ in 200 AGeV Au-Au collisions [1], finding that the suppression pattern of both pions and protons is very similar in the 6+ GeV momentum region. This is to some degree surprising, as the AKK fragmentation functions [2] (which roughly describe the STAR data on proton production in p-p collisions [3]) indicate that $\sim 80\%$ of $p + \bar{p}$ production comes from gluon jets whereas only $\sim 40\%$ of pion production is gluon-driven. This is not so in the older KKP set of fragmentation functions [4] which consequently underpredict proton production at STAR. However, gluons as colour octet states are expected to undergo stronger energy loss in medium than quarks, thus the dominance of gluon jets in hard proton production should map into a stronger suppression of $p + \bar{p}$ production as compared to $\pi^+ + \pi^-$ production. It becomes thus a quantitative question in the energy loss model to check how large this difference in suppression should be and if present data are able to resolve it. A framework to study energy loss within full expansion dynamics of soft matter has been presented in [5] and extended to a 3-d hydrodynamical model [6] in [7]. We utilize this framework to address the above question. We stress that given the scenario describing pionic R_{AA} , the change to study the suppression of protons is only the trivial change of the fragmentation function, no further parameters are adjusted. Furthermore, since the AKK fragmentation function overpredicts $p + \bar{p}$ production in the high p_T region in p-p collisions (and the fragmentation function of $q(\bar{q}) \rightarrow p(\bar{p})$ is well constrained), thus overemphasizing the role of gluons, we stress that our results are an upper limit for the expected difference between proton and pion suppression.

Let us now discuss the treatment of partons propagating

through the medium: our calculation follows the BDMPS formalism for radiative energy loss [8] using quenching weights as introduced by Salgado and Wiedemann [9, 10]. The probability density $P(x_0, y_0)$ for finding a hard vertex at the transverse position $\mathbf{r}_0 = (x_0, y_0)$ and impact parameter $\mathbf{b}$ is given by the product of the nuclear profile functions as

$$P(x_0, y_0) = \frac{T_A(\mathbf{r}_0 + \mathbf{b}/2)T_A(\mathbf{r}_0 - \mathbf{b}/2)}{T_{AA}(\mathbf{b})}, \quad (1)$$

where the thickness function is given in terms of Woods-Saxon the nuclear density $\rho_A(\mathbf{r}, z)$ as $T_A(\mathbf{r}) = \int dz \rho_A(\mathbf{r}, z)$. If we call the angle between outgoing parton and the reaction plane ϕ , the path of a given parton through the medium $\xi(\tau)$ is specified by $(\mathbf{r}_0, \phi)$ and we can compute the energy loss probability $P(\Delta E)_{path}$ for this path. We do this by evaluating the line integrals

$$\omega_c(\mathbf{r}_0, \phi) = \int_0^\infty d\xi \xi \hat{q}(\xi) \quad \text{and} \quad \langle \hat{q}L \rangle(\mathbf{r}_0, \phi) = \int_0^\infty d\xi \hat{q}(\xi) \quad (2)$$

along the path where we assume the relation

$$\hat{q}(\xi) = K \cdot 2 \cdot \epsilon^{3/4}(\xi)(\cosh \rho - \sinh \rho \cos \alpha) \quad (3)$$

between the local transport coefficient $\hat{q}(\xi)$ (specifying the quenching power of the medium), the energy density ϵ and the local flow rapidity ρ with angle α between flow and parton trajectory [11, 12]. Here ω_c is the characteristic gluon frequency, setting the scale of the energy loss probability distribution, and $\langle \hat{q}L \rangle$ is a measure of the path-length weighted by the local quenching power. We view the parameter K as a tool to account for the uncertainty in the selection of α_s and possible non-perturbative effects increasing the quenching power of the medium (see discussion in [5]) and adjust it such that pionic R_{AA} for central Au-Au collisions is described.

Using the numerical results of [9], we obtain $P(\Delta E; \omega_c, R)_{path}$ for ω_c and $R = 2\omega_c^2 / \langle \hat{q}L \rangle$ as a function of jet production vertex and the angle ϕ .

*Electronic address: trenk@phys.jyu.fi

†Electronic address: kari.eskola@phys.jyu.fi




FIG. 1: R_{AA} for pions as measured by PHENIX [13], scaled R_{CP} for pions measured by STAR [1] (in good agreement with the PHENIX R_{AA}), scaled R_{CP} for protons measured by STAR [1] and calculated R_{AA} for pions and protons, dashed lines indicate the regime where recombination is expected to become an important effect [14].

From the energy loss distribution given a single path, we can define the averaged energy loss probability distribution $P(\Delta E)_{TAA}$ as

$$\langle P(\Delta E) \rangle_{TAA} = \frac{1}{2\pi} \int_0^{2\pi} d\phi \int_{-\infty}^{\infty} dx_0 \int_{-\infty}^{\infty} dy_0 P(x_0, y_0) P(\Delta E)_{path}. \quad (4)$$

The energy loss probability $P(\Delta E)_{path}$ is derived in the limit of infinite parton energy [9]. In order to account for the finite energy of the partons we truncate $\langle P(\Delta E) \rangle_{TAA}$ at $\Delta E = E_{jet}$ and add $\delta(\Delta - E_{jet}) \int_{E_{jet}}^{\infty} d\epsilon P(\epsilon)$ to the truncated distribution to ensure proper normalization. The physics meaning of this correction is that we consider all partons as absorbed whose energy loss is formally larger than their initial energy. We calculate the momentum spectrum of hard partons in leading order perturbative QCD (LO pQCD) (explicit expressions are given in [5] and references therein). The medium-modified perturbative production of hadrons can then be computed from the expression

$$d\sigma_{med}^{AA \rightarrow h+X} = \sum_f d\sigma_{vac}^{AA \rightarrow f+X} \otimes \langle P(\Delta E) \rangle_{TAA} \otimes D_{f \rightarrow h}^{vac}(z, \mu_F^2) \quad (5)$$

with $D_{f \rightarrow h}^{vac}(z, \mu_F^2)$ the fragmentation function with momentum fraction z at scale μ_F^2 [2], and from this we compute the nuclear modification factor R_{AA} as

$$R_{AA}(p_T, y) = \frac{dN_{AA}^h/dp_T dy}{T_{AA}(\mathbf{b}) d\sigma^{pp}/dp_T dy}. \quad (6)$$

In Fig. 1 we compare with the STAR data. Since our model, making use of thermodynamical quantities to describe energy loss induced by the soft medium is not suited to describe 60 – 80% peripheral collisions where a dilute, predominantly hadronic, medium is expected




FIG. 2: Calculated R_{AA} for quarks and gluons and protons/antiprotons originating from quarks and gluons using the AKK fragmentation functions as well as the final result averaged over the relative contributions of quarks and gluons in the pQCD calculation.

to exist, we calculate R_{AA} and scale the experimental data. We show that the scaled pionic R_{CP} as obtained by STAR [1] is in perfect agreement with the pionic R_{AA} as obtained by PHENIX [13]. This gives us confidence that the comparison of protonic R_{AA} with scaled R_{CP} is not grossly unreasonable. As apparent from the results, the similarity of the pionic and protonic suppression is expected, the difference between pion and proton R_{AA} is less than the error on the pion R_{AA} over a large kinematic range.

Let us briefly discuss the origin of this outcome: While for an infinite energy parton the average energy loss scales linearly with the coupling strength to the medium (and hence with the color charge), R_{AA} for finite p_T cannot exhibit a linear scaling downward for the simple reason that it is bounded from below by zero. In other words, once the medium quenching is strong enough such that essentially all partons from a spacetime region are absorbed, increasing the quenching power of this region further cannot alter R_{AA} any more and consequently observed hadrons in such a situation are emitted from the medium surface (cf. the discussion in [15]). In the present calculation, we likewise observe this saturation effect (albeit somewhat weaker, cf. [5]), predominantly for gluons.

In Fig. 2 we show R_{AA} for quarks and gluons as well as R_{AA} for protons produced exclusively from quarks and gluons. Since the fraction of protons produced from quarks is small when using the AKK fragmentation functions, the final result for protons is very close to the gluon R_{AA} at all accessible p_T (whereas the pion R_{AA} is between the two partonic curves, showing the transition from a gluon-dominated parton spectrum at low p_T to a quark-dominated spectrum at higher p_T). The gluonic R_{AA} is clearly in a saturated regime and we checked that the spread between quark and gluon R_{AA} is reduced by a factor two as compared to a medium with less quenching

power in which R_{AA} for pions is close to 0.5. It is also apparent that while the functional form of the fragmentation function has some influence on the results at lower p_T , R_{AA} can mainly be understood from the partonic suppression and the p_T dependent ratio of quarks and gluons contributing to the production of a given hadron. Thus, the similarity of R_{AA} for protons and pions in this calculation has to be attributed to the fact that gluon energy loss appears in the saturated regime.

Once again, we stress that we consider the calculation to be an upper limit for the expected difference. For the most part, the STAR data on proton suppression are in a regime where hydrodynamics or recombination [14] is

the dominant contribution to hadron production, in the momentum region above 6 GeV where fragmentation is expected to dominate hadronization, the calculation is in agreement with the data.

Acknowledgments

We thank Jan Rak and Hannu Paukkunen for valuable comments and discussions. This work was financially supported by the Academy of Finland, Project 115262.

-
- [1] B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. Lett. **97** (2006) 152301.
 - [2] S. Albino, B. A. Kniehl and G. Kramer, Nucl. Phys. B **725** (2005) 181.
 - [3] J. Adams *et al.* [STAR Collaboration], Phys. Lett. B **637** (2006) 161.
 - [4] B. A. Kniehl, G. Kramer and B. Potter, Nucl. Phys. B **582**, (2000) 514.
 - [5] T. Renk and K. J. Eskola, hep-ph/0610059.
 - [6] C. Nonaka and S. A. Bass, Phys. Rev. C **75**, 014902 (2007).
 - [7] T. Renk, J. Ruppert, C. Nonaka and S. A. Bass, nucl-th/0611027.
 - [8] R. Baier *et al.*, Nucl. Phys. B **484**, 265 (1997).
 - [9] C. A. Salgado and U. A. Wiedemann, Phys. Rev. D **68**, 014008 (2003).
 - [10] C. A. Salgado and U. A. Wiedemann, Phys. Rev. Lett. **89**, 092303 (2002).
 - [11] R. Baier, A. H. Mueller and D. Schiff, nucl-th/0612068.
 - [12] H. Liu, K. Rajagopal and U. A. Wiedemann, hep-ph/0612168.
 - [13] M. Shimomura [PHENIX Collaboration], nucl-ex/0510023.
 - [14] R. J. Fries *et al.*, Phys. Rev. C **68** (2003) 044902.
 - [15] A. Dainese, C. Loizides and G. Paic, Eur. Phys. J. C **38**, (2005) 461.